

Measure	Description
Prevention & Screening	
Breast Cancer	The percentage of women 40–69 years of age who had a mammogram to screen for breast cancer.
Cervical Cancer	The percentage of women 21–64 years of age who received one or more Pap tests to screen for cervical cancer.
Colorectal Cancer	The percentage of adults 50-80 years of age who had appropriate screening for colorectal cancer (4 criteria applicable).
Chlamydia Screening	The percentage of women 16–25 years of age who were identified as sexually active & who had at least one test for Chlamydia during the measurement year.
Diabetes	The percentage of patients 18-75 years with diabetes (type 1 and type 2):
Eye Exam (Retinal) Performed	Who received a retinal or dilated eye exam by an eye care professional (Ophthalmologist or Optometrist) in the measurement year, or a <i>negative</i> retinal exam (no evidence of retinopathy) by an eye care professional in the year prior to the measurement year.
HgA1C Testing	Who received a HbA1c test performed during the measurement year as identified by claim/encounter or e-laboratory data.
LDL-C Screening	Who received a LDL-C test performed during the measurement year as identified by claim/encounter or e-laboratory data.
Medical Attention for Nephropathy	Who received a nephropathy screening test or evidence of nephropathy as document through electronic data.
Heart Disease	The percentage of patients 18 years of age or older:
Persistence of Beta-Blocker Treatment after a Heart Attack (PBH)	During the measurement year who were hospitalized and discharged alive from 7/1/my-1 through 6/30/my with a diagnosis of AMI and who received persistent beta blocker treatment for six months after discharge. Note: must have pharmacy benefit and continued enrollment through 180 days.

Cholesterol Management for Patients with Cardiovascular Conditions	The percentage of patients 18-75 years who were discharged alive for an AMI, CABG, or PCI from January 1- November 1 of the prior year to the measurement year, or who had a diagnosis of IVD during measurement year and the year prior to the measurement year and who had LDL – C screen and LDL – C control performed.
Measure	Description
Asthma	
Use of appropriate Medications for People with Asthma	The percentage of members 5–56 years of age during the measurement year who were identified as having persistent asthma and who were appropriately prescribed medication during the measurement year.
Pediatrics	
Appropriate Testing for Children with Pharyngitis (CWP)	The percentage of children 2–18 years of age, who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test for the episode. A higher rate denotes better performance.
Appropriate Treatment for Children with Upper Respiratory Infection	The percentage of children 3 months–18 years of age who were given a diagnosis of URI and were not dispensed an antibiotic prescription.
Adults	
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	The percentage of patients 18-64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.
Annual Monitoring For Patients On Persistent Medications	The percentage of patients 18 years of age or older who received at least 180-day supply of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year.