



Wellmark Advantage Health Plan, Inc. is an independent licensee of the Blue Cross and Blue Shield Association.

Medical benefit drug policies are a source for Wellmark Advantage Health Plan medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and therefore subject to change.

Effective Date: 08/04/2022

Pulmonary Arterial Hypertension Products

- Flolan® (epoprostenol)
- Remodulin® (treprostinil)
- Tyvaso® (treprostinil)
- Uptravi® (selexipag)
- Veletri® (epoprostenol)
- Ventavis® (iloprost)

HCPCS: Flolan/Veletri - J1325; Remodulin - J3285; Tyvaso - J7686; Uptravi – J3490; Ventavis - Q4074

Policy:

Requests must be supported by submission of chart notes and patient specific documentation.

- A. Coverage of the requested drug is provided when all the following are met:
 - a. FDA approved indication
 - b. FDA approved age
 - c. If the requested drug is listed below, the member must meet the additional criteria listed:

Drug(s)	Criteria
Tyvaso	<ul style="list-style-type: none"> • For pulmonary hypertension associated with interstitial lung disease (PH-ILD; World Health Organization (WHO) Group 3) - no further criteria are required. • Pulmonary Arterial Hypertension (PAH) WHO Group 1 - trial and failure, intolerance or contraindication to all of the following <ul style="list-style-type: none"> ○ Generic sildenafil or tadalafil • Generic ambrisentan or bosentan

This policy and any information contained herein is the property of Wellmark Advantage Health Plan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and Wellmark Advantage Health Plan employees for the purpose of coverage determinations.

Drug(s)	Criteria
Uptravi injection	<ul style="list-style-type: none"> • Trial and failure, intolerance or contraindication to all of the following: <ul style="list-style-type: none"> ○ Generic sildenafil or tadalafil ○ Generic ambrisentan or bosentan ○ Adempas • Currently stable on oral Uptravi therapy • Will be used as short-term bridging therapy in those patients temporarily unable to take oral therapy
Ventavis	<ul style="list-style-type: none"> • Trial and failure, intolerance or contraindication to all of the following: <ul style="list-style-type: none"> ○ Generic sildenafil or tadalafil • Generic ambrisentan or bosentan

- d. Trial and failure, contraindication, or intolerance to the preferred drugs as listed in Wellmark Advantage Health Plan's prior authorization and step therapy documents and/or Wellmark Advantage Health Plan's utilization management medical drug list

B. Quantity Limitations, Authorization Period and Renewal Criteria

- a. Quantity Limits: Align with FDA recommended dosing
- b. Authorization Period: One year at a time unless specified below
 - i. Uptravi injection: 1 month
- c. Renewal Criteria: Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit unless specified below.
 - i. Uptravi injection: Not applicable as no further authorization will be provided

***Note: Coverage may differ for Medicare Part B members based on any applicable criteria outlined in Local Coverage Determinations (LCD) or National Coverage Determinations (NCD) as determined by Center for Medicare and Medicaid Services (CMS). See the CMS website at <http://www.cms.hhs.gov/>. Determination of coverage of Part B drugs is based on medically accepted indications which have supported citations included or approved for inclusion determined by CMS approved compendia.

Background Information

- FDA approved indications for medications covered in this policy:
 - Epoprostenol: A prostacyclin vasodilator indicated for the treatment of PAH (WHO Group I) to improve exercise capacity. Studies establishing effectiveness included predominantly patients with New York Heart Association (NYHA) Functional Class III-IV symptoms and etiologies of idiopathic or heritable PAH or PAH associated with connective tissue diseases.
 - Treprostinil: A prostacyclin vasodilator indicated for:
 - Treatment of PAH (WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%).

This policy and any information contained herein is the property of Wellmark Advantage Health Plan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and Wellmark Advantage Health Plan employees for the purpose of coverage determinations.

- Patients who require transition from epoprostenol, to reduce the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition.
- Uptravi: A prostacyclin receptor agonist indicated for the treatment of PAH (WHO Group 1) to delay disease progression and reduce the risk of hospitalization for PAH.
- Ventavis: A prostacyclin mimetic indicated for the treatment of PAH (WHO Group 1) to improve a composite endpoint consisting of exercise tolerance, symptoms (NYHA Class), and lack of deterioration. Studies establishing effectiveness included predominately patients with NYHA Functional Class III-IV symptoms and etiologies of idiopathic or heritable PAH (65%) or PAH associated with connective tissue diseases (23%).
- CHEST 2019 guideline supported therapies for WHO Functional Class II and III include: sildenafil, tadalafil, ambrisentan, bosentan, macitentan, riociguat, treprostinil inhalation, and iloprost inhalation
- Additional CHEST 2019 guideline supported therapies for WHO Functional Class III and IV include: epoprostinil IV and treprostinil IV/SQ, and in those who cannot use IV/SQ, treprostinil inhalation and iloprost inhalation.
- PDE5Is and ERAs are clinically effective guideline supported therapies that have wide utility per CHEST guideline recommendations and are cost effective as they are available as generics.
- Opsumit (macitentan) has no clinical advantages over ambrisentan and bosentan at this time, as all three ERAs are now FDA approved to prevent disease progression and clinical worsening. Ambrisentan and bosentan are more cost effective therapies as they are available generically.
- CHEST 2019 guidelines suggest initial monotherapy with ERA, PDE5I or Adempas (riociguat) for treatment of WHO functional class II or III; and suggest adding a second class if inadequate response to initial monotherapy.
- CHEST 2019 guidelines suggest adding a third drug class in WHO Class III or IV with deterioration.
- CHEST 2019 guidelines suggest inhaled prostacyclins (Tyvaso (treprostinil), Ventavis (iloprost)) for WHO Class III after failure of one or two classes of oral agents.
- Cross benefit opportunities may exist to step on drugs on the medical benefit for prostacyclin drugs per guidelines, as SQ/IV therapy is preferred to inhalation WHO Class III and WHO Class IV. Prostacyclin SQ/IV medications may be more cost-effective, however require more complex administration. Further investigation is warranted.
- Uptravi (selexipag) and Orenitram ER (treprostinil tablets) were found to have insufficient clinical evidence to support a guideline recommendation per the CHEST 2019 guidelines. Utilization of guideline supported agents is promoted by step therapy.
- Per package labeling, in Uptravi (selexipag) clinical trials at baseline, 80% of patients were being treated with a stable dose of an ERAs (15%), a PDE5Is (32%), or both (33%).
- Per FDA label, Adempas (riociguat) should not be used in combination with a PDE5I.
- Pulmonary hypertension (PH) with interstitial lung disease and pulmonary fibrosis can be treated with IV or inhaled treprostinil. PDE5Is have shown little or no benefit in these patients. ERAs, specifically ambrisentan, have been shown to be ineffective and associated with adverse effects in patients with PH while bosentan and macitentan are ineffective in idiopathic pulmonary fibrosis (IPF) but have not been tested in IPF associated PH.

- Upravi's prospective, multicenter, open-label, single-sequence, cross-over, Phase III study showed the safety, tolerability and pharmacokinetics of temporarily switching between oral Upravi and Upravi IV in 20 patients. Patients who were stable on oral Upravi switched to IV Upravi for three infusions including the morning and evening dose on Day 1, and morning dose of Day 2 before switching back to the oral formulation in the evening of Day 2.

References:

1. Micromedex. DrugDex. Accessed 9/14/2020 www.micromedex.com
2. Revatio Prescribing Information. New York, NY: Pfizer Labs. April 2020.
3. Adcirca Prescribing Information. Indianapolis, IN: Eli Lilly. August 2020
4. Tracleer Prescribing Information. South San Francisco, CA: Actelion Pharmaceuticals. July 2021
5. Letairis Prescribing Information. Foster City, CA: Gilead Sciences. August 2019
6. Opsumit Prescribing Information. South San Francisco, CA: Actelion Pharmaceuticals. October 2021
7. Upravi Prescribing Information. South San Francisco, CA: Actelion Pharmaceuticals October 2021
8. Adempas Prescribing Information. Whippany, NJ: Bayer HealthCare. September 2021
9. Orenitram ER Prescribing Information. Research Triangle Park, NC; United Therapeutics Corp. May 2021
10. Tyvaso Prescribing Information. Research Triangle Park, NC; United Therapeutics Corp. March 2021
11. Tyvaso DPI Prescribing Information. Research Triangle Park, NC; United Therapeutics Corp. May 2022
12. Ventavis Prescribing Information. Titusville, NJ: Actelion Pharmaceuticals. March 2022
13. Tadalafil (tadalafil) [Prescribing Information]. Farmville, NC; CMP Pharma, Inc. June 2022
14. Klinger, et al. PAH WHO Group 1 CHEST Guidelines 2019. CHEST, 2019. [https://journal.chestnet.org/article/S0012-3692\(19\)30002-9/abstract](https://journal.chestnet.org/article/S0012-3692(19)30002-9/abstract)
15. FDA Drug Safety Communication: FDA recommends against use of Revatio in children with pulmonary hypertension. August 30, 2012. [cited September 11, 2012]; Available from: <http://www.fda.gov/Drugs/DrugSafety/ucm317123.htm>
16. The Regence Group. Preliminary Medication Review: Cardiovascular: sildenafil suspension. September 2012.
17. The Regence Group. Medical Policy for Adcirca and Cialis. July 2012.
18. McLaughlin VV, Badesch DB, Farber HW, et al for the American College of Cardiology Foundation Task Force. ACCF/AHA 2009 Expert Consensus Document on Pulmonary Hypertension: A Report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association. *Circulation* 2009; 119:2250-94.
19. Michelakis E et al. Oral sildenafil is an effective and specific pulmonary vasodilator in patients with pulmonary arterial hypertension. *Circulation* 2002; 105:2398.
20. Ghofrani HA et al. Sildenafil for treatment of lung fibrosis and pulmonary hypertension: a randomized controlled trial. *Lancet* 2002; 360:895-900.
21. Ghofrani HA et al. Sildenafil for long-term treatment of nonoperable chronic thromboembolic pulmonary hypertension. *Am J Respir Crit Care Med* 2003; 167:1139
22. Jackson G et al. Sildenafil for primary pulmonary hypertension: short and long-term symptomatic benefit. *Int J Clin Pract* 2002; 56(5):397-8.
23. Galiè N, Brundage BH, Ghofrani HA, Oudiz RJ, Simonneau G, Safdar Z, Shapiro S, White RJ, Chan M, Beardsworth A, Frumkin L, Barst RJ; Pulmonary Arterial Hypertension and Response to Tadalafil (PHIRST) Study Group. Tadalafil therapy for pulmonary arterial hypertension. *Circulation*. 2009 Jun 9; 119(22):2894-903. Epub 2009 May 26. PubMed PMID: 19470885.
24. Marion, DW. Pulmonary hypertension. In: UpToDate, Basow, DS (Ed), UpToDate, Waltham, MA, 2013.
25. Therapeutic Class ReviewSM. Pulmonary Arterial Hypertension (PAH): Updated for treprostinil (TyvasoTM), tadalafil (AdcircaTM), sildenafil (Revatio[®]), treprostinil (Remodulin[®]) iloprost (Ventavis[®]). The Regence Group. July 2010.
26. Badesch DB, Abman SH, Simonneau G, et al. Medical therapy for pulmonary arterial hypertension: updated ACCP evidence-based clinical practice guidelines. *Chest* 2007; 131:1917.
27. BCBSA Technology Evaluation Center. TEC Specialty Pharmacy Report #15-2013 and #16-2013: Macitentan (Opsumit) and Riociguat (Adempas). October 2013.

28. Galie N, Corris PA, Frost A, Girgis RE, Granton J, Jing ZC, et al. Updated treatment algorithm of pulmonary arterial hypertension. J Am Coll Cardiol 2013;62:D60-72.
29. McLaughlin VV, Gaine, SP, Howard LS, Leuchte HH, Mathier MA, Mehta S, et al. Treatment goals of pulmonary hypertension. J Am Coll Cardio 2013;62:D73-81.
30. Hoeper MM, Bogaard HJ, Condliffe R, Frantz R, Khanna D, Kurzyna M, et al. Definitions and diagnosis of pulmonary hypertension. J Am Coll Cardio 2013;62:D42-50.
31. United Therapeutics Corp. FDA approved Orientram™ (treprostinil) extended-release tablets for the treatment of pulmonary arterial hypertension. Silver Springs, MD: United Therapeutics, 23 December 2013. Web 14 February 2014.

Policy/UM Medical Management System Update History		
#	Date	Change Description
1.1	Effective Date: 08/04/2022	Updated to add Tyvaso DPI formulation and Tadliq
1.0	Effective Date: 01/01/2022	New policy.

** The prescribing information for a drug is subject to change. To ensure you are reading the most current information it is advised that you reference the most updated prescribing information by visiting the drug or manufacturer website or <http://dailymed.nlm.nih.gov/dailymed/index.cfm>.*