



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

## DRUG POLICY

# Radicava (edaravone) Radicava ORS (edaravone)

### NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

### BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

### DESCRIPTION

The indications below including FDA-approved indications and compendial uses are considered covered benefits provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-Approved Indications

Radicava is indicated for the treatment of amyotrophic lateral sclerosis (ALS).

### POLICY

#### Prescriber Specialties

This medication must be prescribed by or in consultation with a neuromuscular specialist.

#### Required Documentation

The following information is necessary to initiate the prior authorization review:

1. Documentation supporting ALSFRS-R score
2. Documentation of respiratory function

The following information is necessary to review requests for continuation of therapy:

1. Documentation supporting clinical benefit from Radicava therapy
2. Documentation supporting patient's ventilator or tracheostomy status

### Criteria for Initial Approval

Authorization of 6 months may be granted for treatment of ALS when all of the following criteria are met:

1. Diagnosis of definite or probable ALS
2. Duration of ALS is 2 years or less from symptom onset
3. Scores of 2 points or better on each individual item of the ALS Functional Rating Scale – Revised [ALSFRS-R] confirmed by medical records
4. Normal respiratory function retained (defined as percent predicted forced vital capacity (FVC) values of 80%) confirmed by medical records
5. Concomitant use of riluzole (at up to maximally indicated doses) unless contraindicated or clinically significant adverse effects are experienced

### Continuation of Therapy

Authorization of 6 months may be granted for members continuing with Radicava therapy when the following criteria are met:

1. Diagnosis of definite or probable ALS
2. There is a clinical benefit from Radicava therapy (e.g., the ability to perform most activities of daily living and not experiencing rapid disease progression while on therapy) as confirmed by medical records
3. Not dependent on invasive ventilation or tracheostomy confirmed by medical records

Radicava is considered **not medically necessary** for members who do not meet the criteria set forth above.

### Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

### Quantity Limits

<b>Medication</b>	<b>Standard Limit</b>	<b>FDA-recommended Dosing</b>
Radicava 30 mg/100 mL per IV bag	600 mg (20 bags) per 28 days	<u>Initial Treatment Cycle:</u> 60 mg/day for 14 days, followed by a 14-day drug-free period  <u>Subsequent Treatment Cycles:</u> 60 mg/day for 10 days out of 14 days, followed by a 14- day drug-free period
Radicava ORS Starter Kit 735 mg/35 mL (105 mg/5 mL dose)	Initial Treatment Cycle: 70mL (2 bottles) per 28 days	<u>Initial Treatment Cycle:</u> 5 mL (105mg)/day for 14 days, followed by a 14-day drug-free period  <u>Subsequent Treatment Cycles:</u>
Radicava ORS Kit 1050 mg/50 mL (105 mg/5 mL dose)	50mL (1 bottle) per 28 days	5 mL (105mg)/day for 10 days out of 14 days, followed by a 14-day drug-free period

## **APPENDIX**

ALS functional rating scale (revised) (ALSFRS-R) is a commonly used functional rating system for persons with ALS. The ALSFRS-R scale consists of 12 questions that evaluate the fine motor, gross motor, bulbar, and respiratory function of patients with ALS (speech, salivation, swallowing, handwriting, cutting food,

dressing/hygiene, turning in bed, walking, climbing stairs, dyspnea, orthopnea, and respiratory insufficiency). Each item is scored from 0 to 4, with higher scores representing greater functional ability.

- Item 1: Speech
  - 4 Normal speech processes
  - 3 Detectable speech disturbance
  - 2 Intelligible with repeating
  - 1 Speech combined with nonvocal communication
  - 0 Loss of useful speech
- Item 2: Salivation
  - 4 Normal
  - 3 Slight but definite excess of saliva in mouth; may have nighttime drooling
  - 2 Moderately excessive saliva; may have minimal drooling
  - 1 Marked excess of saliva with some drooling
  - 0 Marked drooling; requires constant tissue or handkerchief
- Item 3: Swallowing
  - 4 Normal eating habits
  - 3 Early eating problems — occasional choking
  - 2 Dietary consistency changes
  - 1 Needs supplemental tube feeding
  - 0 NPO (exclusively parenteral or enteral feeding)
- Item 4: Handwriting
  - 4 Normal
  - 3 Slow or sloppy: all words are legible
  - 2 Not all words are legible
  - 1 Able to grip pen but unable to write
  - 0 Unable to grip pen
- Item 5a: Cutting food and handling utensils (patients without gastrostomy -> use item 5b if >50% is through g-tube)
  - 4 Normal
  - 3 Somewhat slow and clumsy, but no help needed
  - 2 Can cut most foods, although clumsy and slow; some help needed
  - 1 Food must be cut by someone, but can still feed slowly
  - 0 Needs to be fed
- Item 5b: Cutting food and handling utensils (alternate scale for patients with gastrostomy)
  - 4 Normal
  - 3 Clumsy but able to perform all manipulations independently
  - 2 Some help needed with closures and fasteners
  - 1 Provides minimal assistance to caregiver
  - 0 Unable to perform any aspect of task
- Item 6: Dressing and hygiene
  - 4 Normal function
  - 3 Independent and complete self-care with effort or decreased efficiency
  - 2 Intermittent assistance or substitute methods
  - 1 Needs attendant for self-care
  - 0 Total dependence
- Item 7: Turning in bed and adjusting bed clothes
  - 4 Normal
  - 3 Somewhat slow and clumsy, but no help needed
  - 2 Can turn alone or adjust sheets, but with great difficulty
  - 1 Can initiate, but not turn or adjust sheets alone
  - 0 Helpless
- Item 8: Walking
  - 4 Normal

- 3 Early ambulation difficulties
- 2 Walks with assistance
- 1 Nonambulatory functional movement
- 0 No purposeful leg movement
- Item 9: Climbing stairs
  - 4 Normal
  - 3 Slow
  - 2 Mild unsteadiness or fatigue
  - 1 Needs assistance
  - 0 Cannot do
- Item 10: Dyspnea
  - 4 None
  - 3 Occurs when walking
  - 2 Occurs with one or more of the following: eating, bathing, dressing (ADL)
  - 1 Occurs at rest, difficulty breathing when either sitting or lying
  - 0 Significant difficulty, considering using mechanical respiratory support
- Item 11: Orthopnea
  - 4 None
  - 3 Some difficulty sleeping at night due to shortness of breath, does not routinely use more than two pillows
  - 2 Needs extra pillows in order to sleep (more than two)
  - 1 Can only sleep sitting up
  - 0 Unable to sleep
- Item 12: Respiratory insufficiency
  - 4 None
  - 3 Intermittent use of BiPAP
  - 2 Continuous use of BiPAP during the night
  - 1 Continuous use of BiPAP during the night and day
  - 0 Invasive mechanical ventilation by intubation or tracheostomy

## PROCEDURES AND BILLING CODES

**To report provider services, use appropriate CPT\* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.**

- C9493 – injection, edaravone, 1mg (deleted 12-31-2018)
- J1301 – Radicava, Injection, edaravone, 1mg (effective 1-1-2019)
- J8499 – Prescription drug, oral, non chemotherapeutic, nos (Radicava ORS (edaravone))

## REFERENCES

- Radicava and Radicava ORS [package insert]. Jersey City, NJ: MT Pharma America, Inc.; March 2022.
- EFNS Task Force on Diagnosis and Management of Amyotrophic Lateral Sclerosis; Andersen PM, et al. EFNS guidelines on the Clinical Management of Amyotrophic Lateral Sclerosis (MALS) – revised report of an EFNS task force. *Eur J Neurol.* 2012;19(3):360-75.
- Cedarbaum JM, Stambler N, Malta E, et al. The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. BDNF ALS Study Group (Phase III). *J Neurol Sci.* 1999;169(1-2):13-21.
- Abe K, Itoyama Y, Sobue G, et al. Confirmatory double-blind, parallel-group, placebo-controlled study of efficacy and safety of edaravone (MCI-186) in amyotrophic lateral sclerosis patients. *Amyotroph Lateral Scler Frontotemporal Degener.* 2014; 15(7-8):610-617.
- Brooks BR, Miller RG, Swash M, Munsat TL; World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic

lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord.* 2000; 1(5):293-299.

- Chiò A, Logroscino G, Traynor BJ, et al. Global epidemiology of amyotrophic lateral sclerosis: a systematic review of the published literature. *Neuroepidemiology.* 2013; 41(2):118-130.
- Douglass CP, Kandler RH, Shaw PJ, McDermott CJ. An evaluation of neurophysiological criteria used in the diagnosis of motor neuron disease. *J Neurol Neurosurg Psychiatry.* 2010; 81(6):646-649.
- Writing Group; Edaravone (MCI-186) ALS 19 Study Group. Safety and efficacy of edaravone in well defined patients with amyotrophic lateral sclerosis: a randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* 2017 May 15.
- Yoshino H, Kimura A. Investigation of the therapeutic effects of edaravone, a free radical scavenger, on amyotrophic lateral sclerosis (Phase II study). *Amyotroph Lateral Scler.* 2006; 7(4):241-245.
- Luo L et al. Efficacy and safety of edaravone in treatment of amyotrophic lateral sclerosis-a systematic review and meta-analysis. *Neurol Sci.* 2019;40(2):235-241.
- Shefner J et al. Long-term edaravone efficacy in amyotrophic lateral sclerosis: Post-hoc analyses of Study 19 (MCI186-19). *Muscle Nerve.* 2020;61(2):218-221.

\*Some content reprinted from CVSHealth

## POLICY HISTORY

**Policy #:** 05.02.25

**Reviewed:** July 2022

**Revised:** July 2022

**Current Effective Date:** October 1, 2022