



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

Simponi Aria (golimumab injection for intravenous use)

NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

DESCRIPTION

The intent of the Simponi Aria drug policy is to ensure appropriate selection of patients for therapy based on product labeling, clinical guidelines and clinical studies. The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

1. Adult patients with moderately to severely active rheumatoid arthritis (RA) in combination with methotrexate
2. Active psoriatic arthritis (PsA) in patients 2 years of age and older
3. Adult patients with active ankylosing spondylitis (AS)
4. Active polyarticular Juvenile Idiopathic Arthritis (pJIA) in patients 2 years of age and older

Compendial Uses

Oligoarticular juvenile idiopathic arthritis

POLICY

Documentation

- A. Rheumatoid arthritis (RA)
 1. For initial requests:
 - i. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

- ii. Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).
 - 2. For continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- B. Psoriatic arthritis: For continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- C. Ankylosing spondylitis (AS) and articular juvenile idiopathic arthritis:
- 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable). If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

Criteria for Initial Approval

A. Moderately to severely active rheumatoid arthritis (RA)

- 1. Authorization of 12 months may be granted for members who have previously received a biologic or targeted synthetic DMARD (e.g. Rinvoq, Xeljanz) indicated for moderately to severely active rheumatoid arthritis. The requested medication must be prescribed in combination with methotrexate unless the member has a clinical reason not to use methotrexate (see Appendix A).
- 2. Authorization of 12 months may be granted for treatment of moderately to severely active RA when all of the following criteria are met:
 - i. Member meets either of the following criteria:
 - a. Member has been tested for either of the following biomarkers and the test was positive:
 - 1. Rheumatoid factor (RF)
 - 2. Anti-cyclic citrullinated peptide (anti-CCP)
 - b. Member has been tested for ALL of the following biomarkers:
 - 1. RF
 - 2. Anti-CCP
 - 3. C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
 - ii. Member is prescribed the requested medication in combination with methotrexate or has a clinical reason not to use methotrexate.
 - iii. Member meets either of the following criteria:
 - a. Member has experienced an inadequate response to at least a 3-month trial of methotrexate despite adequate dosing (i.e., titrated to at least 15 mg/week).
 - b. Member has an intolerance or contraindication to methotrexate (See Appendix A).

B. Active psoriatic arthritis (PsA)

- 1. Authorization of 12 months may be granted for treatment of active psoriatic arthritis (PsA).

C. Active ankylosing spondylitis (AS)

- 1. Authorization of 12 months may be granted for members who have previously received a biologic indicated for active ankylosing spondylitis.
- 2. Authorization of 12 months may be granted for treatment of active ankylosing spondylitis when any of the following criteria is met:
 - a. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - b. Member has an intolerance or contraindication to two or more NSAIDs.

D. Active articular juvenile idiopathic arthritis

1. Authorization of 12 months may be granted for members who have previously received a biologic or targeted synthetic DMARD indicated for active articular juvenile idiopathic arthritis
2. Authorization of 12 months may be granted for treatment of active articular juvenile idiopathic arthritis when any of the following criteria is met:
 - a). The member has had an inadequate response to methotrexate or another non-biologic DMARD administered at an adequate dose and duration.
 - b). The member has risk factors (See Appendix C) and the member also meets one of the following:
 - a. High-risk joints are involved (e.g., cervical spine, wrist, or hip).
 - b. High disease activity.
 - c. Are judged to be at high risk for disabling joint disease.

Continuation of Therapy

A. Moderately to severely active rheumatoid arthritis (RA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement

C. Active ankylosing spondylitis (AS)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active ankylosing spondylitis and who achieve or maintain a positive clinical response with the requested medication as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Functional status
2. Total spinal pain
3. Inflammation (e.g., morning stiffness)

D. Active articular juvenile idiopathic arthritis

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active articular juvenile idiopathic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
2. Number of joints with limitation of movement
3. Functional ability

Other

For all indications: Member has had a documented negative TB test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for

persons who are naïve to biologic DMARDs or targeted synthetic DMARDs associated with an increased risk of TB.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic DMARD or targeted synthetic DMARD.

Simponi Aria is considered **not medically necessary** for members who do not meet the criteria set forth above.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Appendix

Appendix A: Examples of Contraindications to Methotrexate

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia)
4. Elevated liver transaminases
5. History of intolerance or adverse event
6. Hypersensitivity
7. Interstitial pneumonitis or clinically significant pulmonary fibrosis
8. Myelodysplasia
9. Pregnancy or currently planning pregnancy
10. Renal impairment
11. Significant drug interaction

Appendix B: Examples of Contraindications to the Use of NSAIDs

1. Allergic-type reaction following aspirin or other NSAID administration
2. Asthma
3. Gastrointestinal bleeding
4. History of intolerance or adverse event
5. Significant drug interaction
6. Urticaria

Appendix C: Risk factors for Articular Juvenile Idiopathic Arthritis

1. Positive rheumatoid factor
2. Positive anti-cyclic citrullinated peptide antibodies
3. Pre-existing joint damage

PROCEDURES AND BILLING CODES

To report provider services, use appropriate CPT* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.

- J1602 Injection, golimumab, 1 mg, for intravenous use

REFERENCES

- Simponi Aria [package insert]. Horsham, PA: Janssen Biotech, Inc.; September 2020.
- Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis*. 2020;79(6):685-699. doi:10.1136/annrheumdis-2019-216655.
- Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2016;68(1):1-26.
- Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum*. 2008;59(6):762-784.
- Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
- Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2016;75(3):499-510.
- Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol*. 2019;71(10):1599-1613. doi:10.1002/art.41042.
- Tuberculosis (TB). TB risk factors. Centers for Disease Control and Prevention. Retrieved on 12 August 2020 from: <https://www.cdc.gov/tb/topic/basics/risk.htm>.
- Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol*. 2019;71(1):5-32. doi:10.1002/art.40726.
- Ringold S, Angeles-Han S, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *American College of Rheumatology*. 2019;1-18.
- Aletaha D, Neogi T, Silman, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum*. 2010;62(9):2569-81.
- Clinical Consult: CVS Caremark Clinical Programs Review. Focus on Rheumatology Clinical Programs. March 2021.
- Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
- Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care Res*. 2021;0:1-16.

*Some content reprinted from CVSHealth

POLICY HISTORY

Policy #: 05.02.13

Reviewed: April 2022

Revised: May 2021

Current Effective Date: July 1, 2022