



Wellmark Blue Cross and Blue Shield is an independent licensee of the Blue Cross and Blue Shield Association.

DRUG POLICY

Tremfya (guselkumab)

NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

DESCRIPTION

The intent of the Tremfya drug policy is to ensure appropriate selection of patients for therapy based on product labeling, clinical guidelines and clinical studies.

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

1. Treatment of adult patients with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy
2. Treatment of adult patients with active psoriatic arthritis

POLICY

Required Documentation

Submission of the following information is necessary to initiate the prior authorization review:

A) Plaque psoriasis

1. Initial requests:
 - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected.
 - ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.

B) Psoriatic arthritis:

1. For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

Criteria for Initial Approval

A. Moderate to severe plaque psoriasis

1. Authorization of 12 months may be granted for members who have previously received Otezla or a biologic indicated for the treatment of moderate to severe plaque psoriasis.
2. Authorization of 12 months may be granted for treatment of moderate to severe plaque psoriasis for members when all of the following criteria are met:
 - a) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - b) Member meets any of the following criteria:
 - i.) Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine or acitretin.
 - ii.) Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine and acitretin (see Appendix B).
 - iii.) Member has severe psoriasis that warrants a biologic DMARD as first-line therapy (i.e. at least 10% of the body surface area (BSA) or crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected).

B. Active psoriatic arthritis (PsA)

1. Authorization of 12 months may be granted for treatment of active psoriatic arthritis.

Continuation of Therapy

A. Moderate to severe plaque psoriasis (PsO)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active psoriatic arthritis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement

Other

For all indications: Member has had a documented negative TB test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic DMARDs or targeted synthetic DMARDs associated with an increased risk of TB.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic DMARD or targeted synthetic DMARD.

Tremfya is considered **not medically necessary** for members who do not meet the criteria set forth above.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Quantity Limits

Trade Name	Generic Name	Quantity Limit	FDA-recommended dosing
Tremfya®	guselkumab	Initiation of therapy: 2 syringes per first 28 days Maintenance: 1 syringe per 56 days	<ul style="list-style-type: none"> • Loading doses: 100 mg at week 0 and 4 • Maintenance dose: 100 mg every 8 weeks thereafter

Appendix

Appendix A: Clinical reasons to avoid TNF-inhibitors

1. History of demyelinating disorder
2. History of congestive heart failure
3. History of hepatitis B infection
4. Autoantibody formation/lupus-like syndrome
5. Risk of lymphoma

Appendix B: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine or Acitretin

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Cannot be used due to risk of treatment-related toxicity
4. Drug interaction
5. Pregnancy or currently planning pregnancy
6. Significant comorbidity prohibits use of systemic agents (examples include liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

PROCEDURES AND BILLING CODES

To report provider services, use appropriate CPT* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.

- N/A

REFERENCES

- Tremfya [package insert]. Horsham, PA: Janssen Biotech, Inc.; July 2020.
- Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. J Am Acad Dermatol. 2009;61:451-485.

- Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
- Reich K, Armstrong, AW, Foley P, et al. Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the treatment of patients with moderate to severe psoriasis with randomized withdrawal and retreatment: Results from the phase III, double-blind, placebo- and active comparator–controlled VOYAGE 2 trial. *Am J Clin Dermatol*. 2017;76(3):418-431.
- Blauvelt A, Papp KA, Griffiths, CEM, et al. Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the continuous treatment of patients with moderate to severe psoriasis: Results from the phase III, double-blinded, placebo- and active comparator-controlled VOYAGE 1 trial. *Am J Clin Dermatol*. 2017;76(3):405-417.
- Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80(4):1029-1072.
- Tuberculosis (TB). TB risk factors. Centers for Disease Control and Prevention. Retrieved on 15 November 2021 from: <https://www.cdc.gov/tb/topic/basics/risk.htm>.

*Some content reprinted from CVSHealth

POLICY HISTORY

Policy #: 05.02.20

Reviewed: April 2022

Revised: April 2022

Current Effective Date: July 21, 2022