



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

## DRUG POLICY

# Sporanox and Onmel

### NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

### BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

### DESCRIPTION

The intent of the Sporanox and Onmel prior authorization program is to ensure appropriate selection of patients based on product labeling and/or clinical guidelines and/or clinical studies, and to encourage the use of more cost-effective agents prior to the use of Sporanox or Onmel for the treatment of onychomycosis..

The Food and Drug Administration (FDA) has approved the use of **Sporanox® (itraconazole) capsules** for treatment of the following fungal infections:

- Blastomycosis, pulmonary and extrapulmonary
- Histoplasmosis, including chronic cavitary pulmonary disease and disseminated, non-meningeal histoplasmosis, and
- Aspergillosis, pulmonary and extrapulmonary, in patients who are intolerant of or who are refractory to amphotericin B therapy.

**Sporanox® (itraconazole) capsules** are also approved for the following onychomycosis diagnoses:

- Onychomycosis of the toenail, with or without fingernail involvement, due to dermatophytes (tinea unguium)
- Onychomycosis of the fingernail due to dermatophytes (tinea unguium).

Compendial uses for **Sporanox® (itraconazole) capsules**

- Coccidioidomycosis
- Cryptococcosis
- Microsporidiosis
- Penicilliosis

- Sporotrichosis

The FDA has approved the use of **Sporanox® (itraconazole) oral solution** for the treatment of oropharyngeal and esophageal candidiasis.

Compendial uses for **Sporanox® (itraconazole) solution**

- Blastomycosis
- Histoplasmosis
- Aspergillosis
- Coccidioidomycosis
- Cryptococcosis
- Microsporidiosis
- Penicilliosis
- Sporotrichosis

**Onmel (itraconazole) tablets** are FDA approved for the treatment of onychomycosis of the toenail caused by *Trichophyton rubrum* or *Trichophyton mentagrophytes*.

**POLICY**

- I. **Sporanox capsule (brand and generic)** therapy may be considered **medically necessary** for the treatment of systemic fungal infections or for prophylaxis of invasive fungal infections in post-transplant patients

**Approval** will be for up to **12 months**.

- II. **Sporanox capsule (brand and generic)** may be considered **medically necessary** for the treatment of onychomycosis when the following criteria are met:
  - Diagnosis has been confirmed by appropriate laboratory testing (e.g., potassium hydroxide [KOH] preparation, fungal culture, or nail biopsy)  
AND
  - Treatment is considered medically necessary (e.g., not for cosmetic purposes only)  
AND
  - Patient has experienced an inadequate response, adverse event, intolerance, or contraindication to terbinafine therapy

**Approval** will be for **12 weeks**. (Only 1 approval allowed in a 12 month period.)

- III. **Onmel** may be considered **medically necessary** for the treatment of onychomycosis when the following criteria are met:
  - Diagnosis has been confirmed by appropriate laboratory testing (e.g., potassium hydroxide [KOH] preparation, fungal culture, or nail biopsy)  
AND
  - Treatment is considered medically necessary (e.g., not for cosmetic purposes only)  
AND
  - Patient has experienced an inadequate response, adverse event, intolerance, or contraindication to terbinafine therapy  
AND
  - The patient is unable to take generic Sporanox due to an allergy, intolerance, or contraindication to the excipients

**Approval will be for 12 weeks.** (Only 1 approval allowed in a 12 month period.)

IV. ***Sporanox oral solution*** may be considered **medically necessary** when the following criteria are met:

- Those who have a diagnosis of oropharyngeal and/or esophageal candidiasis AND have experienced an inadequate response, adverse event, intolerance, or contraindication to fluconazole therapy  
OR
- Those with a diagnosis of a systemic fungal infection who cannot physically swallow itraconazole capsules or unable to achieve therapeutic levels with itraconazole capsules  
OR
- Patient has a life-threatening or serious infection

**Approval will be for up to 12 months.**

V. **The aforementioned drugs are considered not medically necessary for patients who do not meet the criteria set forth above.**

#### Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

### PROCEDURES AND BILLING CODES

***To report provider services, use appropriate CPT\* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.***

- Code(s), if applicable.

### REFERENCES

- Sporanox® (itraconazole) capsules [package insert]. Titusville, NJ: Ortho-McNeil/Janssen Pharmaceuticals Inc, April 2012.
- Sporanox® (itraconazole) oral solution [package insert]. Raritan, NJ: Centocor Ortho Biotech Products, L.P., March 2012.
- Onmel™ (itraconazole) tablets [package insert]. Research Triangle Park, NC: Stiefel Laboratories, Inc, April 2010.
- Itraconazole. *Drug Facts and Comparisons*. Facts & Comparisons® eAnswers [online]. 2013. Available from Wolters Kluwer Health, Inc. Accessed January 14, 2013.
- De Berker D. Fungal nail disease. *N Engl J Med* 2009; 360:2108-2116.
- Roberts DT, Taylor WD, Boyle J. Guidelines for treatment of onychomycosis. *British Journal of Dermatology*. 2003; 148: 402-410
- Gupta AK, Drummond-Main C, Cooper EA, et al. Systematic review of nondermatophyte mold onychomycosis: diagnosis, clinical types, epidemiology and treatment. *J Am Acad Dermatol* 2012; 66: 494:502.
- Gupta AK, et al. Prevalence and Epidemiology of toenail onychomycosis in diabetic subjects: a multicentre survey. *Br J Dermatology*. 1998; 139:665-671.
- Elewski BE. Large-scale epidemiological study of the causal agents of onychomycosis: mycological findings from the multicenter onychomycosis study of terbinafine. *Arch Dermatology*. 1997; 133:1317-1318.

- Jaffe, R. Onychomycosis: recognition, diagnosis, and management. *Arch Fam Med*. 1998; 7:587-592.
- Rodgers P, Bassler M MD. Treating Onychomycosis. *Am Fam Physician*. 2001; 63: 663-72, 677-8.
- Scher RK. Onychomycosis: a significant medical disorder. *J Am Acad Dermatol*. 1996; 35(3 pt 2): S2-5.
- Goldenstein AO, Goldenstein BG. Onychomycosis. In: *UpToDate*, Rose BD (Ed.), UpToDate. Waltham, MA, 2012.
- Kauffman CA. Treatment of Oropharyngeal and Esophageal Candidiasis. In: *UpToDate*, Rose BD (Ed.), UpToDate. Waltham, MA, 2012.
- De Berker D. Fungal nail disease. *N Engl J Med* 2009;360:2108-2116.
- Tosti A. Onychomycosis. Medscape [online serial]. Updated July 28, 201. Available at: <http://emedicine.medscape.com/article/1105828-overview#a0156>. Accessed September 17, 2014.
- Elewski BE, Rich P, Pollak R, et al. Efinaconazole 10% solution in the treatment of toenail onychomycosis: two phase III multicenter, randomized, double-blind studies. *J Am Acad Dermatol*. 2013;68(4):600-608.
- Westerberg DP, Voyack MJ. Onychomycosis: current trends in diagnosis and treatment. *Am Fam Physician*. 2013;88(11):762-770.
- Thomas J, Jacobson GA, Markowicz CK, Peterson GM, Burnet H, Sharpe C. Toenail onychomycosis: an important global disease burden. *J Clin Pharm Ther*. 2010;35(5):497-519.
- Crawford F, Young P, Godfrey C, et al. Oral treatments for toenail onychomycosis. A systematic review. *Arch Dermatol* 2002;138:811-816.
- Gupta AK, Ryder JE, Johnson AM. Cumulative meta-analysis of systemic antifungal agents for the treatment of onychomycosis. *Br J Dermatol*. 2004;150(3):537-544.
- Favero ML, Bonetti AF, Domingos EL et al. Oral antifungal therapies for toenail onychomycosis: a systematic review and meta-analysis toenail mycosis: network meta-analysis. *J Dermatolog Treat*. 2020;1-10. doi: 10.1080/09546634.2020.1729336. Online ahead of print. Accessed September 11, 2020.
- Sporanox Oral Capsules (itraconazole). CVS Caremark Criteria. Ref # 280-A. Accessed September 14, 2020.
- Bhatia A, Kanish B, Badyal DK et al. Efficacy of oral terbinafine versus itraconazole in treatment of dermatophytic infection of skin – A prospective, randomized comparative study. *Indian J Pharmacol*. 2019;51(2):116-119. Accessed September 21, 2020.
- Itraconazole Oral. Facts & Comparisons. Updated September 17, 2020. Accessed September 21, 2020.
- 5.01.18 Itraconazole. BlueCross BlueShield Federal Employee Program. Updated December 6, 2019. Accessed September 21, 2020.

## POLICY HISTORY

**Policy #:** 05.01.76

**Policy Creation:** April 2006

**Reviewed:** October 2020

**Revised:** February 2019

**Current Effective Date:** April 7, 2019